European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA


La neurotoxicité liée à l'anesthésie : effet de la chirurgie (et de sa cause)

Congrès commun ADARPEF/ESPA, Bordeaux, France 2 Avril 2016

Tom G. Hansen, MD, PhD,
Department of Anaesthesia & Intensive Care - University Hospital
& University of Southern Denmark DENMARK
Email: tomghansen@dadlnet.dk




OUH


Hans Christian Andersen
The Emperor's New Clothes


Trollhättan
Falköping
Hirtshals
Hjorring
Göteborg
Frederikshavn
Boras
Thisted
Ålborg
Varberg
Struer
Skive
Halmstad
Denmark
Ringkobing
Grena
Herning
Langå
Århus
Skjern
Hässleholm
Esbjerg
Copenhagen
Kastrup
Malmo
Odense
Nyborg
Tonder
Trelleborg
Niebüll
Flensburg
Nykobing
Husum
Rødby
Heide
Kiel
Rendsburg
Puttgarden
Stralsund
Binz
Norddeich
Cuxhaven
Neumünster
Rostock
Emden
Wilhelmshaven
Hamburg
Schwerin
Zussow
Oldenburg
Leer
Bremen
Newbrandenburg
Ludwigslust
Lüneburg
Neustrelitz
Uelzen
Wittenberge
Celle
Wolfsburg
Angermunde
Osnabrück



ESPA

Sujets de l'exposé

COMMENT LA CHIRURGIE PEUT-ELLE INFLUENCER LE DEVELOPPEMENT NEURO-DEVELOPEMENTAL DE L'ENFANT


- TRAUMATISME , p ex STRESS, INFLAMMATION
- DOULEUR
- LE DIAGNOSTIC / LES PATHOLOGIES ASSOCIEES
- (ANESTHESIE)

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Blockade of NMDA Receptors and Apoptotic Neurodegeneration in the Developing Brain

Chrysanthy Ikonomidou,* Friederike Bosch, Michael Miksa, Petra Bittigau, Jessica Vöckler, Krikor Dikranian, Tanya I. Tenkova, Vanya Stefovska, Lechoslaw Turski, John W. Olney



Science 1999;283:70-74

Early Exposure to Common Anesthetic Agents Causes Widespread Neurodegeneration in the Developing Rat Brain and Persistent Learning Deficits

Vesna Jevtovic-Todorovic,[1] Richard E. Hartman,[2] Yukitoshi Izumi,[1] Nicholas D. Benshoff,[1] Krikor Dikranian,[3] Charles F. Zorumski,[1] John W. Olney,[3] and David F. Wozniak[3]
[1]Department of Anesthesiology, University of Virginia Health System, Charlottesville, Virginia 22908, and Departments of [2]Neurology and [3]Psychiatry, Washington University School of Medicine, St. Louis, Missouri 63110

Etudes sur des modèles animaux

Repeated administration of ketamine may lead

Poten … **in**

comb … **the infant**

mous

[1]Chainll … [1]Haihui Wang,

A. [1]Joann …

*Divisi

Long-t … **ngle**

episod … **lockade**

G Andr…

Alicia S…

Elizabet…

Rat Brain

Chrysanth … and Persistent Learning Deficits

Petra

Tanya I.



Vesna Jevtovic-Todorovic,[1] Richard E. Hartman,[2] Yukitoshi Izumi,[3] Nicholas D. Benshoff,[3] Krikor Dikranian,[3] Charles F. Zorumski,[3] John W. Olney,[3] and David F. Wozniak[3]

[1]Department of Anesthesiology, University of Virginia Health System, Charlottesville, Virginia 22908, and Departments of [2]Neurology and [3]Psychiatry, Washington University School of Medicine, St. Louis, Missouri 63110

SINGLE-DOSE KETAMINE ADMINISTRATION INDUCES APOPTOSIS IN NEONATAL MOUSE BRAIN

Michael Rudin,[1] Ron Ben-Abraham,[2] Vered Gazit,[3] Yevgeney Tendler,[3], Vadim Tashlykov,[3], and Yeshayahu Katz

[1]Dept. of Anesthesiology, HaEmek Medical Center, Afula, Israel; [2]Dept. of Anesthesiology and Critical Care, Tel Aviv Sourasky Medical Center and Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; [3]Laboratory of Anesthesia, Pain and Neural Research, The Bruce Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel

Etudes sur des cohortes humaines

Attention-Deficit/Hyperactivity Disorder After Early Exposure to Procedures Requiring General Anesthesia

Juraj Sprung, MD, PhD; Randall P. Flick, MD, MPH; Slavica K. Katusic, MD; Robert C. Colligan, PhD; William J. Barbaresi, MD; Katarina Bojanić, MD; Tasha L. Welch, MD; Michael D. Olson, PA-C; Andrew C. Hanson, BS; Darrell R. Schroeder, MS; Robert T. Wilder, MD, PhD; and David O. Warner, MD

Abstract

Objective: To study the association between exposure to procedures performed under general anesthesia before age 2 years and development of attention-deficit/hyperactivity disorder (ADHD).
Patients and Methods: Study patients included all children born between January 1, 1976, and December 31, 1982, in Rochester, MN, who remained in Rochester after age 5. Cases of ADHD diagnosed before age 19 years were identified by applying stringent research criteria. Cox proportional hazards regression assessed exposure to procedures requiring general anesthesia (none, 1, 2 or more) as a predictor of ADHD using a stratified analysis with strata based on a propensity score including comorbid health conditions.
Results: Among the 5357 children analyzed, 341 ADHD cases were identified (estimated cumulative incidence, 7.6%; 95% confidence interval [CI], 6.8%-8.4%). For children with no postnatal exposure to procedures requiring anesthesia before the age of 2 years, the cumulative incidence of ADHD at age 19 years was 7.3% (95% CI, 6.5%-8.1%). For single and 2 or more exposures, the estimates were 10.7% (95% CI, 6.8%-14.4%) and 17.9% (95% CI, 7.2%-27.4%), respectively. After adjusting for gestational age, sex, birth weight, and comorbid health conditions, exposure to multiple (hazard ratio, 1.95; 95% CI, 1.03-3.71), but not single (hazard ratio,1.18; 95% CI, 0.79-1.77), procedures requiring general anesthesia was associated with an increased risk for ADHD.
Conclusion: Children repeatedly exposed to procedures requiring general anesthesia before age 2 years are at increased risk for the later development of ADHD even after adjusting for comorbidities.

© 2012 Mayo Foundation for Medical Education and Research Mayo Clin Proc. 2012;87(2):120-129

Cognitive and Behavioral Outcomes After Early Exposure to Anesthesia and Surgery

AUTHORS: Randall P. Flick, MD, MPH,* Slavica K. Katusic, MD,† Robert C. Colligan, PhD,‡ Robert T. Wilder, MD, PhD,* Robert G. Voigt, MD,§ Michael D. Olson, PhD,* Juraj Sprung, MD, PhD,* Amy L. Weaver, MS,† Darrell R. Schroeder, MS,† and David O. Warner, MD*

*Department of Anesthesiology, †Division of Epidemiology, Department of Health Sciences Research, ‡Department of Psychiatry and Psychology, †Division of Biomedical Statistics and Informatics, Department of Health Sciences Research, and §Department of Pediatrics and Adolescent Medicine, and *Mayo Clinic Mayo Medical School, College of Medicine, Mayo Clinic, Rochester, Minnesota

WHAT'S KNOWN ON THIS SUBJECT: Exposure to virtually all anesthetic drugs has been shown to cause neurodegeneration in young animals. Studies of learning and cognition in children exposed to anesthesia and surgery have been few, have relied on single outcome measures, and have not controlled for comorbidity.

WHAT THIS STUDY ADDS: In this study of children exposed to anesthesia/surgery before the age of 2, multiple group and individual measures of learning and behavior are examined by using a matched design with adjustment for comorbidity using 2

Anesthesiology 2009; 110:796–804 Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Early Exposure to Anesthesia and Learning Disabilities in a Population-based Birth Cohort

Robert T. Wilder, M.D., Ph.D., Randall P. Flick, M.D., M.P.H.,† Juraj Sprung, M.D., Ph.D.,‡ Slavica K. Katusic, M.D.,§ William J. Barbaresi, M.D.,‖ Christopher Mickelson, M.D.,# Stephen J. Gleich, M.D.,** Darrell R. Schroeder, M.S.,†† Amy L. Weaver, M.S.,†† David O. Warner, M.D.‡*

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Anesthesia for Cesarean Delivery and Learning Disabilities in a Population-based Birth Cohort

Juraj Sprung, M.D., Ph.D., Randall P. Flick, M.D., M.P.H.,† Robert T. Wilder, M.D., Ph.D.,‡ Slavica K. Katusic, M.D.,§ Tasha L. Pike, M.S.,‖ Mariella Dingli, M.D.,# Stephen J. Gleich, M.D.,** Darrell R. Schroeder, M.S.,†† William J. Barbaresi, M.D.,‡‡ Andrew C. Hanson, B.S.,§§ David O. Warner, M.D.**



Etudes
sur des cohortes humaines

Are Anesthesia and Surgery durin
with Altered Academic Performan

Robert I. Block, Ph.D.,* Joss J. Thomas, M.D.,† Emine O. E
Karolie K. Kimble, R.N., B.A.,§ Michael M. Todd, M.D.‖

Anesth Analg. 2011 Nov;113(5):1143-51. Epub 2011 Mar 17.

Early childhood exposure to anesthesia and risk of developmental and behavioral disorders in a sibling birth cohort.

DiMaggio C, Sun LS, Li G.
Department of Anesthesiology, Columbia University, 622 West 168 St., New York, NY 10032, USA. cjd11@columbia.edu

Anesthesiology 2009; 1

Behavio
of First

Cor J. Kalkman,
Marcel Bruens, F

A

Long-term Differences in Language and Cognitive Function After Childhood Exposure to Anesthesia

WHAT'S KNOWN ON THIS SUBJECT: Immature animals exposed to anesthetics display apoptotic neurodegeneration and long-term cognitive deficiencies. In children, studies of cognitive deficits associated with anesthesia exposure have yielded mixed results. No studies to date have used directly administered neuropsychological assessments as outcome measures.

WHAT THIS STUDY ADDS: This study examines the association between exposure to anesthesia in children under age 3 and deficits at age 10 by using a battery of directly administered neuropsychological assessments, with deficits found in language and abstract reasoning associated with exposure.

AUTHORS: Caleb Ing, MD,[a] Charles DiMaggio, PhD,[a,b,c] Andrew Whitehouse, PhD,[d] Mary K. Hegarty, MBBS, FANZCA,[e] Joanne Brady, MS,[a,b,c] Britta S. von Ungern-Sternberg, ProfPhD,[e,f] Andrew Davidson, MD,[g] Alastair J.J. Wood, MD,[h] Guohua Li, MD,[a,b,c] and Lena S. Sun, MD[a,i]

Departments of [a]Anesthesiology, [b]Epidemiology, and [i]Pediatrics, Columbia University College of Physicians and Surgeons, New York, New York; [c]Mailman School of Public Health, New York, New York; [d]Centre for Child Health Research and Neurocognitive Development Unit, School of Psychology, and [f]School of Medicine and Pharmacology, The University of Western Australia, Perth, Australia [e]Department of Anaesthesia and Pain Management, Princess Margaret Hospital for Children, Perth, Australia; [g]Department of Anaesthesia, Murdoch Childrens Research Institute & Royal Children's Hospital, Melbourne, Australia; and [h]Department of Medicine, Weill Cornell College of Medicine, and Symphony Capital LLC, New York, New York

Abstra
used anesthetic agents may have serious neurotoxic effects on the developing brain. The purpose of this study was to assess the association between surgery for hernia repair and the risk of behavioral and developmental disorders in young children. We performed a retrospective cohort analysis of children who were enrollees of the New York State Medicaid program. Our

Key Words: anesthesia, surgery, behavior, development
(*J Neurosurg Anesthesiol* 2009;21:286–291)

ESPA

Limites des études de cohortes

- Petit nombre de patients
- Etudes monocentriques
- Différents types de chirurgie
- Large éventail d'âges
- Pas de détails sur l'anesthésie
- Cohortes protégées (région) ou associées à un centre hospitalier
- Critère évalué (difficultés d'apprentissage, hyperactivité, autisme, grades scolaires, tests neuropsychologiques)
- Facteurs confondants connus et inconnus
- Problèmes de migration familiale

ESPA

Critères évalués dans les études de cohorte

- Retard de développement (école gardienne)
- Difficultés d'apprentissage (école primaire)
- Problèmes du comportement (retard mental, autisme/hyperactivité et troubles de l'attention)
- Problèmes psychiatriques
- Troubles du langage
- Succès des études ou QI?
- Tests cognitifs individuels ou de groupe

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

BMJ

BMJ 2014;348:g3387 doi: 10.1136/bmj.g3387 (Published 30 May 2014)

ANALYSIS

Is animal research sufficiently evidence based to be a cornerstone of biomedical research?


Public acceptance of the use of animals in biomedical research is conditional on it producing benefits for humans. **Pandora Pound** and **Michael Bracken** argue that the benefits remain unproved and may divert funds from research that is more relevant to doctors and their patients

Pandora Pound *medical sociologist*[1], Michael B Bracken *Susan Dwight Bliss professor of epidemiology*[2]

[1]Bath, UK; [2]Yale University Schools of Public Health and Medicine, New Haven CT, USA

* La réalisation, la synthèse et la publication de nombreuses études animales restent inadéquates
* La situation actuelle n'est pas éthique car des animaux et des hommes sont utilisés pour des recherches qui ne produiront pas de résultats fiables
* Il n'y a pas d'évidence suffisante de l'utilité des études animales
* Davantage de rigueur et de responsabilité sont nécessaires pour une meilleure gestion des fonds publics

ESPA


Pathologie
anomalie génétique
malformations
prématurité
sepsis
Développement
Chirurgie
stress hormonal
inflammation
dépressioncardiorespiratoire
température
Anesthésie
effets protecteurs
vs toxiques
autres agents “toxiques”
p ex oxygène

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Pediatric Anesthesia

Pediatric Anesthesia ISSN 1155-5645

REVIEW ARTICLE

Anesthesia-related neurotoxicity and the developing animal brain is not a significant problem in children

Tom G. Hansen[1,2]

1 Department of Anesthesiology and Intensive Care –Pediatric Section, Odense University Hospital, Odense C, Denmark
2 Clinical Institute – Anesthesiology, University of Southern Denmark, Odense C, Denmark

La neurotoxicité liée à l'anesthésie
sur le cerveau en développement de l'animal
n'est pas un problème majeur pour l'enfant

Taille de l'échantillon ?

Anesthesiology 2009; 110:805–12 Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Behavior and Development in Children and Age at the Time of First Anesthetic Exposure

Cor J. Kalkman, M.D., Ph.D., Linda Peelen, M.Sc.,† Karel G. Moons, Ph.D.,* Mona Veenhuizen, M.D.,‡ Marcel Bruens, R.N.,§ Gerben Sinnema, Ph.D.,‖ Tom P. de Jong, M.D., Ph.D.#*

Test: child behavioural checklist

N= 243 (0-6 ans) urologie: pas de différence

Calcul a posteriori de la taille de l'échantillon

Enfants > 2 ans => n > 6.000

Nourrissons < 6 mois => n > 2.200

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Age, taille de l'échantillon, chirurgie ?

PERIOPERATIVE MEDICINE

Academic Performance in Adolescence after Inguinal Hernia Repair in Infancy

A Nationwide Cohort Study

Tom G. Hansen, M.D., Ph.D.,* Jacob K. Pedersen, M.Sc.,† Steen W. Henneberg, M.D., Ph.D.,‡
Dorthe A. Pedersen, M.Sc.,† Jeffrey C. Murray, M.D.,§ Neil S. Morton, M.D.,‖
Kaare Christensen, M.D., Ph.D., D.M.Sc.#

Résultats scolaires chez des adolescents qui ont bénéficié d'une cure de hernie inguinale quand ils étaient nourrissons

ESPA

Critères évalués dans les études de cohorte danoises

1. Résultats scolaires en 9ème niveau de scolarité

(0-13; moyenne = 8, SD=1)

Tests et évaluation par l'enseignant (Danois, langues étrangères, Maths, Sciences biologiques & sociales)

2. en cas d'échec

Ajustement en fonction de: p.ex.: sexe, âge, poids de naissance, malformations congénitales, âge et niveau de scolarité des parents

ESPA

Statistiques danoises- les registres

- La banque des données démographiques
- Le registre national des hôpitaux
- La banque des données du marché de l'emploi
- Le registre des examens scolaires obligatoires et de leurs résultats

Europea
www.euro

ESPA

Table 1. Characteristics of Early Anesthetic Exposure *versus* Control Group: Denmark, 1986–1990 Birth Cohort

Characteristic	Exposure Group (n = 2,689)	Control Group (n = 14,575)
Death	34 (1.3)	183 (1.3)
1–365 d after birth	6 (0.2)	109 (0.7)
>365 d after birth	28 (1.0)	74 (0.5)
Migration	109 (4.1)	752 (5.2)
Study base	2,547 (94.7)	13,640 (93.6)
Congenital malformation	316 (12.4)	624 (4.6)
Deceased	8 (23.5)	39 (21.3)
Migrant	9 (8.3)	33 (4.4)
Birth weight	2,541 (99.8)	13,616 (99.8)
<1,500 g	152 (6.0)	70 (0.5)
Birth weight, g	3,000 ± 798	3,437 ± 556
≤1,499	25 (16.4)	11 (15.7)
≥1,500	291 (12.2)	610 (4.5)
Sex (boy)	2,266 (89.0)	6,955 (51.0)
Paternal age, yr	30.3 ± 5.4	30.7 ± 5.4
Maternal age, yr	27.8 ± 4.9	28.1 ± 4.8
Paternal education	1.49 ± 1.71	1.68 ± 1.78
Maternal education	1.53 ± 1.66	1.77 ± 1.74
Average test score	7.73 ± 1.09	7.99 ± 1.08
Boy	7.68 ± 1.08	7.83 ± 1.08
Girl	8.08 ± 1.06	8.14 ± 1.06
Average teacher score	7.71 ± 1.11	8.03 ± 1.09
Boy	7.65 ± 1.11	7.80 ± 1.10
Girl	8.20 ± 1.01	8.25 ± 1.04
Test score	2,016 (79.2)	11,850 (86.9)
Boy	1,771 (78.2)	5,843 (84.0)
Girl	245 (87.2)	6,007 (89.9)
Teacher score	2,028 (79.6)	11,893 (87.2)
Boy	1,782 (78.6)	5,876 (84.5)
Girl	246 (87.5)	6,017 (90.0)

All data are presented as No. (%) or mean ± SD unless otherwise specified.

Groupe exposé

contrôle

E-mail: info@eur

garçons filles

hernie

non opérés




Fig. 1. Distribution of ninth grade mean test scores for Danish adolescents (born 1986–1990) who underwent inguinal hernia repair in infancy (n = 2,547) *versus* a control group composed of a randomly selected 5% population sample of age-matched cohorts (n = 14,575).

What We Already Know about This Topic

- Animal studies have raised concerns about the potential for neurocognitive impairment from general anesthesia in human newborns, but clinical studies are inconclusive.

What This Article Tells Us That Is New

Sur un collectif de > 2.500 enfants opérés de hernie inguinale quand ils étaient nourrissons, aucune différence dans les tests scolaires en 9ème niveau après ajustement pour les facteurs confondants connus

Rapport des cotes en faveur d'un échec du test scolaire: 1,13 (95% IC : 0,98-1,31)

Pediatric Anesthesia

Pediatric Anesthesia ISSN 1155-5645

ORIGINAL ARTICLE

Educational outcome in adolescence following pyloric stenosis repair before 3 months of age: a nationwide cohort study

Tom G. Hansen[1,2], Jacob K. Pedersen[3], Steen W. Henneberg[4], Neil S. Morton[5] & Kaare Christensen[3,6,7]

1 Department of Anesthesia & Intensive Care, Odense University Hospital, Odense C, Denmark
2 Clinical Institute – Anesthesiology, University of Southern Denmark, Odense C, Denmark
3 Institute of Public Health – Epidemiology, University of Southern Denmark, Odense C, Denmark
4 Department of Anesthesia, The Juliane Marie Center, Copenhagen University Hospital –Rigshospitalet, Copenhagen, Denmark
5 Department of Anesthesia, Royal Hospital for Sick Children, Glasgow, UK
6 Department of Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense C, Denmark
7 Department of Clinical Genetics, Odense University Hospital, Odense C, Denmark

Résultats scolaires chez des adolescents qui ont bénéficié d'une cure de sténose de pylore avant l'âge de 3 mois

Euro
www

esthesiology

ESPA


Table 1 Characteristics of pyloric stenosis patients and 5% population sample from the 1986–90 birth cohorts in Denmark

	Pyloric stenosis		5%-Population sample	
	n	Percent (%)	*n*	Percent (%)
No of live births	779		14 666	
Deaths (days after birth)	6	0.8	185	1.3
1–365	3	0.4	110	0.8
>365	3	0.4	75	0.5
Migrations	25	3.2	757	5.2
Cong. malf. among deceased[a]	6	100.0	40	21.6
Cong. malf. among migrants[a]	21	84.0	31	4.1
Study base	748	96.0	13 723	93.6
Congenital malformations[a]	674	90.1	595	4.3
Among birth weights −1499	2	33.3	12	14.8
Among birth weights 1500+	671	90.6	581	4.3
Congenital malformations excl. Pyl. Sten.[b]	51	6.8	583	4.4
Among birth weights −1499	0	0.0	12	14.8
Among birth weights 1500+	50	6.7	569	4.3
Known birth weight	747	99.9	13 700	99.8
Birth weight <1500 (g)	6	0.8	81	0.6
Males	603	80.6	7032	51.2
	Mean	sd	Mean	sd
Birth weight (g)	3345	624	3434	560
Maternal age	27.9	4.9	28.0	4.8
Paternal age	30.7	5.6	30.7	5.4
Maternal education	1.57	1.68	1.77	1.74
Paternal education	1.49	1.74	1.67	1.78
Average test score	7.82	1.15	7.99	1.08
Males	7.78	1.14	7.83	1.08
Females	7.98	1.18	8.14	1.06
Average teachers' score	7.81	1.17	8.03	1.09
Males	7.74	1.19	7.80	1.10
Females	8.06	1.06	8.25	1.04
	n	Percent (%)	*n*	Percent (%)
Available test score	586	79.7	11 910	86.8
Males	470	77.9	5898	83.9
Females	126	86.9	6012	89.9
Available teachers' score	605	80.9	11 954	87.1
Males	478	79.3	5932	84.4
Females	127	87.6	6022	90.0

All available individuals included.
[a]Any congenital malformation as expressed by an ICD-8 code of 740–759.99 or an ICD-10 code of Q00.0–Q99.9.
[b]Any congenital malformation other than Pyloric Stenosis, as expressed by an ICD-8 code of 740–759.99 excluding 750.10–750.19 or an ICD-10 code of Q00.0–Q99.9 excluding Q40.0.

Sténose pylore

contrôles



garçons filles

Rapport des cotes en faveur d'un échec du test scolaire:

1,37 (95% IC: 1,11-1,68)

E-mail: i

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Neurodevelopmental outcome at 2 years of age after general anaesthesia and awake-regional anaesthesia in infancy (GAS): an international multicentre, randomised controlled trial

*Andrew J Davidson, Nicola Disma, Jurgen C de Graaff, Davinia E Withington, Liam Dorris, Graham Bell, Robyn Stargatt, David C Bellinger, Tibor Schuster, Sarah J Arnup, Pollyanna Hardy, Rodney W Hunt, Michael J Takagi, Gaia Giribaldi, Penelope L Hartmann, Ida Salvo, Neil S Morton, Britta S von Ungern Sternberg, Bruno Guido Locatelli, Niall Wilton, Anne Lynn, Joss J Thomas, David Polaner, Oliver Bagshaw, Peter Szmuk, Anthony R Absalom, Geoff Frawley, Charles Berde, Gillian D Ormond, Jacki Marmor, Mary Ellen McCann, for the GAS consortium**

www.thelancet.com Vol 387 January 16, 2016

Résultats à 2 ans de l'étude GAS: une anesthésie générale réalisée dans la petite enfance et qui dure moins d'1 heure, n'entraîne pas, à l'âge de 2 ans, de différence cliniquement significative du développement neuro-comportemental par comparaison à une ALR vigile.

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Pediatric Anesthesia
Pediatric Anesthesia ISSN 1155-5645

Pediatric Anesthesia 25 (2015) 186–192

ORIGINAL ARTICLE

Neurosurgical conditions and procedures in infancy are associated with mortality and academic performances in adolescence: a nationwide cohort study

Tom G. Hansen[1,2], Jacob K. Pedersen[3], Steen W. Henneberg[4], Neil S. Morton[5] & Kaare Christensen[3,6,7]

Association entre neurochirurgie chez le nourrisson, mortalité et résultats scolaires à l'adolescence

Test scolaire au 9ème niveau Neurochirugie chez le nourrisson Cohorte danoise naissance 1986-1990	Mortalité (%)	Echec du test (%)	Scores du Test
Toutes neurochirurgies (n=228)	15,8	48,7	7,57
Hydrocéphalie (n=130)	18,5	59,7	7,51
Craniotomie (n=43)	18,6	40,0	7,32
Myélo/encéphalocèles (n=55)	7,3	32,7	7,78
Contrôles (n=14,703)	1,3	13,2	7,99

European Society for Paediatric Anaesthesiology
www.euroespa.com

ESPA

Après cure d'une fente faciale

Cohorte danoise, naissance 1986-1990 (âge 15-16 ans)	Age lors de l'opération	Mortalité (%)	N qui ont passé le test du 9ème niveau (%)	Score du test (SD)
Toutes formes de fentes (n=588)	2m – 22 m	1,5	81,5	7,93 (1,09)
Fente labiale (n=171)	**2m**-6m	1,2	88,4	8,10 (1,04)
Fente palatine (n=195)	14m-**22m**	1,0	72,1	7,85 (1,10)
Fente labiopalatine (n=222)	**2m**-6m 14-**22m**	2,3	84,4	7,83 (1,12)
Contrôles (n=14.677)	-	1,3	86,8	7,99 (1,09)

E-mail: info@euroespa.com | Web: www.euroespa.com

Facteurs compensatoires possibles pour les fentes labiales: meilleure éducation des parents? moins de prématurés? environnement + stimulant? distribution différentes des sexes?

ACTA OTO-LARYNGOLOGICA, 2016
http://dx.doi.org/10.3109/00016489.2016.1146412

Taylor & Francis
Taylor & Francis Group

RESEARCH ARTICLE

School performance in cholesteatoma-operated children in Denmark: a nationwide population-based register-study

B. D. Djurhuus[a], T. G. Hansen[b], J. K. Pedersen[c], C. E. Faber[a] and K. Christensen[c]

[a]Department of ENT Head & Neck Surgery, Odense University Hospital, Odense, Denmark; [b]Department of A
Odense University Hospital, Odense, Denmark; [c]Epidemiology, Biostatistics and Biodemography, Institute of Pu
Denmark, Odense, Denmark

ESPA

Résultats scolaires chez enfants opérés d'un cholestéatome

Table 1. Characteristics of children surgically treated for middle ear cholesteatoma before 15 years of age (cholesteatoma group) and a 5% population sample (control group) from the Danish 1986–1991 birth cohorts.

	Cholesteatoma group		Control group		
	n	%	*n*	%	*p*-value[c]
Total number	702		17 757		
Exclusions					
Congenital malformations[a]	114	16.2	1362	7.7	<0.001
Emigrated before 17 years	16	2.7	774	4.7	0.024
Died before 17 years	1	0.2	121	0.8	0.067
Unknown birth weight	1	0.2	27	0.2	1.000
Different grading scale	18	3.2	346	2.2	0.147
Unknown parental education	3	0.5	21	0.1	0.051
Study base	549	78.2	15 106	85.1	<0.001
Sex					
Males	343	62.5	7627	50.6	<0.001
Females	206	37.5	7479	49.5	<0.001
	Mean	SD	Mean	SD	*p*-value[d]
âge 1ère opération	9.4	3.1	–	–	–
Birth weight (g)	3393	578	3448	587	0.015
Maternal education[b]	2.1	0.8	2.1	0.7	0.217
Paternal education[b]	1.9	0.7	2.0	0.7	0.015
Average grade					
Males	7.7	1.1	7.8	1.1	0.110
Females	8.1	1.1	8.2	1.1	0.140
9th grade marks registered	*N*	%	*n*	%	*p*-value[c]
Males	290	84.6	6708	88.0	0.060
Females	182	88.4	6915	92.5	0.029

Table 3. Results from three different linear regressions adjusting for the same confounders as in Table 2 (not shown).

	Coefficient	95% CI		*p*-value
Mathematics				
Sex				
Male	Ref	–		–
Female	0.12	0.08;	0.16	<0.001
No. cholesteatoma surgeries				
0	Ref	–		–
1	−0.06	−0.19;	0.08	0.413
≥2	−0.02	−0.19;	0.16	0.869
Danish				
Sex				
Male	Ref	–		–
Female	0.61	0.5;	0.65	<0.001
No. cholesteatoma surgeries				
0	Ref	–		–
1	−0.01	−0.12;	0.10	0.876
≥2	−0.12	−0.27;	0.02	0.101
English				
Sex				
Male	Ref	–		–
Female	0.35	0.30;	0.40	<0.001
N de chirurgie pour cholé				
0	Ref	–		–
1	−0.12	−0.29;	0.04	0.148
> 2	−0.26	−0.48;	−0.03	0.024

European Society for Paediatri
www.euroespa.com

Chirurgie majeure non-cardiaque

ESPA

Neurocognition after neonatal surgery?

Neurodevelopmental Outcomes After Neonatal Surgery for Major Noncardiac Anomalies

Lisanne J. Stolwijk, MD,[a,b,c] Petra MA Lemmers, MD, PhD,[b] Marissa Harmsen, BSc,[a] Floris Groenendaal, MD, PhD,[b] Linda S. de Vries, MD, PhD,[b,c] David C. van der Zee, MD, PhD,[a] Manon J.N. Benders, MD, PhD,[b,c] Maud Y.A. van Herwaarden-Lindeboom, MD, PhD[a]

PEDIATRICS Volume 137, number 2, February 2016

Conclusion: Un retard de développement cognitif et moteur a été observé chez 23 % des patients après une chirurgie néonatale majeure non-cardiaque. Une méta-analyse a montré un score moyen inférieur de 0,5 DS au score normatif de la population en bonne santé

Facteurs de risque:
- **poids de naissance**
- **prématurité**
- **anomalies congénitale**
- **anomalie chromosome**
- **durée hospitalisation**
- **durée ventilation**
- **besoins O_2**
- **retard de croissance**
- **niveau éducation des parents**

Chirurgie cardiaque

European Society for Paediatric Anaes
www.euroespa.com

ESPA

Neurocognition after congenital cardiac surge

Neurodevelopmental Outcomes After Cardiac Surgery in Infancy

J. William Gaynor, MD[a], Christian Stopp, MS[b], David Wypij, PhD[b], Dean B. Andropoulos, MD[c], Joseph Atallah, MD, CM, SM, FRCPC[d], Andrew M. Atz, MD[e], John Beca, MD[f], Mary T. Donofrio, MD[g], Kim Duncan, MD[h], Nancy S. Ghanayem, MD[i], Caren S. Goldberg, MD[j], Hedwig Hövels-Gürich, MD[k], Fukiko Ichida, MD[l], Jeffrey P. Jacobs, MD[m], Robert Justo, MD[n], Beatrice Latal, MD[o], Jennifer S. Li, MD[p], William T. Mahle, MD[q], Patrick S. McQuillen, MD[r], Shaji C. Menon, MD[s], Victoria L. Pemberton, RNC, MS, CCRC[t], Nancy A. Pike, RN, PhD[u], Christian Pizarro, MD[v], Lara S. Shekerdemian, MD[w], Anne Synnes, MDCM[x], Ismee Williams, MD[y], David C. Bellinger, PhD[b], Jane W. Newburger, MD, MPH[b], for the International Cardiac Collaborative on Neurodevelopment (ICCON) Investigators

PEDIATRICS Volume 135, number 5, May 2015

A Systematic Review of Motor and Cognitive Outcomes After Early Surgery for Congenital Heart Disease

AUTHORS: Suzanne H. Snookes, BS,[a,b] Julia K. Gunn, MBBS, FRACP,[c,d,e] Bev J. Eldridge, BS, PhD,[a] Susan M. Donath, BS, MA,[d,f,g] Rod W. Hunt, BMBS, MMed, PhD, FRACP, MRCP,[c,d,e] Mary P. Galea, BS, BA, PhD,[b] and Lara Shekerdemian, MD, MRCP, FRACP[c,d,h]

CONCLUSIONS: With this review we identified a limited number of prospective studies that systematically addressed outcome in patients at the highest risk. These studies consistently revealed cognitive and motor delay in children after cardiac surgery during early infancy. Additional investigation is required to ascertain the consequences of such impairment during later childhood and into adult life. *Pediatrics* 2010; 125:e818–e827


Facteurs de risque:
- poids de naissance
- prématurité
- anomalies congénitale
- anomalie chromosome
- complexité malformation cardiaque
- durée séjour réanimation /hospitalisation
- ventilation préopératoire
- score APGAR
- niveau socioéconomique et éducation et des parents
- ethnie
- technique CEC (Hct, hypoth)
- glycémie

European
www.euroes

Resolving Postoperative Neuroinflammation and Cognitive Decline

Niccolò Terrando, PhD,[1] Lars I. Eriksson, MD, PhD,[1,2] Jae Kyu Ryu, PhD,[3]
Ting Yang, MD, PhD,[1] Claudia Monaco, MD, PhD,[4] Marc Feldmann, FMedSci, FAA, FRS,[4]
Malin Jonsson Fagerlund, MD, PhD,[2] Israel F. Charo, MD, PhD,[5]
Katerina Akassoglou, PhD,[3,6] and Mervyn Maze, MB, ChB[1]

ANN NEUROL 2011;70:986–995



Neuroinflammation postopératoire et déclin cognitif

ESPA

Detection of Postoperative Cognitive Decline After Coronary Artery Bypass Graft Surgery is Affected by the Number of Neuropsychological Tests in the Assessment Battery


Matthew S. Lewis, BAppSc Hons, Paul Maruff, PhD, Brendan S. Silbert, MBBS, FANZCA, Lis A. Evered, BS, and David A. Scott, MBBS, FANZCA, PhD

Centre for Anaesthesia and Cognitive Function, Department of Anaesthesia, St. Vincent's Hospital, Victoria Parade, Melbourne, and School of Psychological Science, La Trobe University, Victoria, Australia

(Ann Thorac Surg 2006;81:2097–104)
© 2006 by The Society of Thoracic Surgeons

- **Il n'existe pas encore d'outils fiables pour évaluer les troubles cognitifs post-opératoires (TCPO)**
- **Il existe un nombre croissant de tests pour mesurer TCPO**
- **Il faut améliorer leur sensibilité dans les groupes exposés et contrôles**

European Society for Paedi
www.euroespa.com



PERIOPERATIVE MEDICINE

Cognitive Functioning after Surgery in Middle-aged and Elderly Danish Twins

Unni Dokkedal, M.P.H., Tom G. Hansen, M.D., Ph.D., Lars S. Rasmussen, M.D., Ph.D., D.M.Sc., Jonas Mengel-From, M.Sc., Ph.D., Kaare Christensen, M.D., Ph.D., D.M.Sc.

Etude comparative sur des jumeaux âgés

ABSTRACT

Background: Postoperative cognitive dysfunction is common, but it remains unclear whether there are long-term adverse cognitive effects of surgery combined with anesthesia. The authors examined the association between exposure to surgery and level of cognitive functioning in a sample of 8,503 middle-aged and elderly twins.

Methods: Results from five cognitive tests were compared in twins exposed to surgery, classified as major, minor, hip and knee replacement, or other, with those of a reference group without surgery using linear regression adjusted for sex and age. Genetic and shared environmental confounding was addressed in intrapair analyses of 87 monozygotic and 124 dizygotic same-sexed twin pairs in whom one had a history of major surgery and the other did not.

Results: Statistically significantly lower composite cognitive score was found in twins with at least one major surgery compared with the reference group (mean difference, −0.27; 95% CI, −0.48 to −0.06), corresponding to one tenth of an SD, that is, a negligible effect size. In the intrapair analysis, the surgery-exposed co-twin had the lower cognitive score in 49% (95% CI, 42 to 56%) of the pairs. None of the other groups differed from the reference group except the knee and hip replacement group that tended to have higher cognitive scores (mean difference, 0.35; 95% CI, −0.18 to 0.87).

Conclusion: un antécédent de chirurgie majeure n'est associé qu'à une diminution mineure des capacités cognitives. Le fonctionnement cognitif préopératoire et les comorbidités ont plus d'influence que la chirurgie et l'anesthésie **(Anesthesiology 2016; 124:312-21)**

Pediatric Anesthesia

EDITORIAL

Anesthetists rather than anesthetics are the threat to baby brains

Markus Weiss[1], Bruno Bissonnette[2,3], Thomas Engelhardt[4], Sulpicio Soriano[5]
[1]Department of Anesthesia, University Children's Hospital, Zurich, Switzerland
[2]Department of Anesthesiology and Pain Medicine, Nationwide Children's Hospital, The Ohio State University, Columbus, OH, USA
[3]Department of Anesthesia and Critical Care Medicine, University of Toronto, Toronto, ON, Canada
[4]Department of Anesthesia, Royal Aberdeen Children's Hospital, Aberdeen, UK
[5]Department of Anesthesia, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA
Email: markus.weiss@kispi.uzh.ch

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Infantile Postoperative Encephalopathy: Perioperative Factors as a Cause for Concern

Mary Ellen McCann, A.N.J. Schouten, Nicole Dobija, Carlos Munoz, Lianne Stephenson, Tina Y. Poussaint, C.J. Kalkman, Paul R. Hickey, Linda S. de Vries and Robert C. Tasker

Pediatrics 2014;133;e751; originally published online February 10, 2014;
DOI: 10.1542/peds.2012-0973

ESPA

Mais qu'en est-il de

Hypo-/hypercarbie?
Hypo- /hyperoxie?
Pression artérielle/ de perfusion?
Hypo/hyper-glycémie?
hyponatrémie?
Hypo/hyperthermie?
Formation/compétenc

REVIEW

CURRENT OPINION

Safe Anesthesia For Every Tot – The SAFETOTS initiative

Markus Weiss[a,b], Laszlo Vutskits[c], Tom G. Hansen[d,e], and Thomas Engelhardt[f]

ESPA

Conclusions

La chirurgie et les comorbidité ont beaucoup d'influence sur le développement neurocognitif

Les études animales ne sont pas prédictives des résultats chez l'humain

Attitude différente en Europe par rapport aux USA ?

Assez de publications sans évidence de preuves, s'il vous plaît!

Autre chose ??